

October 2023

Dear Parents and Carers

Is my child too ill for school?

There is wide agreement among health professionals and educational professionals that school attendance is vital to the life chances of children and young people. Being in school improves health, wellbeing and socialisation throughout the course of life. The greatest benefits come from children and young people attending school regularly.

We know it can be tricky deciding whether or not to keep your child off school when they're unwell. We are aware that the COVID-19 pandemic may have caused some parents to feel less confident with assessing whether their child is well enough to be in school so we have laid out some NHS guidance which we hope you will find helpful. It is usually appropriate for parents and carers to send their children to school with mild respiratory illnesses. This would include general cold symptoms: a minor cough, runny nose or sore throat. However, children should not be sent to school if they have a temperature of 38°C or above.

In addition to respiratory illnesses, we are aware that more children may be absent from school due to symptoms of anxiety than before the pandemic. Worry and mild or moderate anxiety, whilst sometimes difficult emotions, can be a normal part of growing up for many children and young people. Being in school can often help alleviate the underlying issues. A prolonged period of absence is likely to heighten a child's anxiety about attending in the future, rather than reduce it.

If you do keep your child at home, it's important to phone the school on the first day to let them know that your child won't be in and give the reason. A call must then be made every subsequent day until the child returns to school. You may receive a call from the school to discuss their absence and to encourage your child's return as soon as possible to reduce lost learning time.

Other illnesses

Please find below the NHS' advice for common childhood illnesses:

Coughs and colds

It's fine to send your child to school with a minor [cough](#) or [common cold](#). But if they have a fever, keep them off school until the fever goes. Encourage your child to throw away any used tissues and to wash their hands regularly.

High temperature

If your child has a [high temperature](#), (38°C and above) keep them off school until it goes away.

Chickenpox

If your child has [chickenpox](#), keep them off school until all the spots have crusted over. This is usually about 5 days after the spots first appeared.

Cold sores

There's no need to keep your child off school if they have a [cold sore](#). Encourage them not to touch the blister or kiss anyone while they have the cold sore, or to share things like cups and towels.

Conjunctivitis

You don't need to keep your child away from school if they have [conjunctivitis](#). Do get advice from your pharmacist. Encourage your child not to rub their eyes and to wash their hands regularly.

COVID-19

If your child has mild symptoms, such as a runny nose, sore throat, or slight cough, and feels well enough, they can go to school. Your child should try to stay at home and avoid contact with other people if they have [symptoms of COVID-19](#) and they either have a high temperature, do not feel well enough to go to school or do their normal activities.

What to do if your child has tested positive

Your child is no longer required to do a COVID-19 rapid lateral flow test if they have symptoms. But if your child has tested positive for COVID-19, they should try to stay at home and avoid contact with other people for 3 days after the day they took the test.

Ear infection

If your child has an [ear infection](#) and a high temperature or severe earache, keep them off school until they're feeling better or their high temperature goes away.

Hand, foot and mouth disease

If your child has [hand, foot and mouth disease](#) but seems well enough to go to school, there's no need to keep them off. Encourage your child to throw away any used tissues straight away and to wash their hands regularly.

Head lice and nits

There's no need to keep your child off school if they have had head lice. You can treat [head lice and nits](#) without seeing a GP.

Impetigo

If your child has [impetigo](#), they'll need treatment from a GP, often with antibiotics. Keep them off school until all the sores have crusted over and healed, or for 48 hours after they start antibiotic treatment. Encourage your child to wash their hands regularly and not to share things like towels and cups with other children at school.

Ringworm

If your child has [ringworm](#), see your pharmacist unless it's on their scalp, in which case you should see a GP. It's fine for your child to go to school once they have started treatment.

Scarlet fever

If your child has [scarlet fever](#), they'll need treatment with antibiotics from a GP. Otherwise they'll be infectious for 2 to 3 weeks. Your child can go back to school 24 hours after starting antibiotics.

Sore throat

You can still send your child to school if they have a [sore throat](#). But if they also have a high temperature, they should stay at home until it goes away. A sore throat and a high temperature can be symptoms of [tonsillitis](#).

Threadworms

You don't need to keep your child off school if they have [threadworms](#). Speak to your pharmacist, who can recommend a treatment.

Vomiting and diarrhoea

Children with [diarrhoea or vomiting](#) should stay away from school until they have not been sick or had diarrhoea for at least 1 day (24 hours).

Kind regards

The Attendance Team